

Patient Information	Specimen Information	Client Information

COMMENTS:

Test Name	In Range	Out Of Range	Reference Range	Lab
BETA 2 GLYCOPROTEIN I AB (IGG, IGA, IGM)				
B2 GLYCOPROTEIN I (IGG)AB	<2.0		<20.0 U/mL	
Value	Interpretation			
< 20.0	Antibody not detected			
> or = 20.0	Antibody detected			
B2 GLYCOPROTEIN I (IGM)AB	<2.0		<20.0 U/mL	
Value	Interpretation			
< 20.0	Antibody not detected			
> or = 20.0	Antibody detected			
B2 GLYCOPROTEIN I (IGA)AB	<2.0		<20.0 U/mL	
Value	Interpretation			
< 20.0	Antibody not detected			
> or = 20.0	Antibody detected			

The antiphospholipid antibody syndrome (APS) is a clinical-pathologic correlation that includes a clinical event (e.g. arterial or venous thrombosis, pregnancy morbidity) and persistent positive antiphospholipid antibodies (IgM, IgG Cardiolipin or b2GPI antibodies greater than the 99th percentile; or a lupus anticoagulant). International consensus guidelines for APS suggest waiting at least 12 weeks before retesting to confirm antibody persistence. The Systemic Lupus International Collaborating Clinics immunological classification criteria for systemic lupus erythematosus (SLE) include testing for isotype IgA, which has yet to be incorporated into APS criteria. Low level antiphospholipid antibodies may sometimes be detected in the setting of infection, drug therapy or aging.

For additional information, please refer to <http://education.questdiagnostics.com/faq/FAQ109> (This link is being provided for informational/educational purposes only.)

PERFORMING SITE:

SPECIMEN: