

Patient Information	Specimen Information	Client Information

COMMENTS:

Test Name	In Range	Out Of Range	Reference Range	Lab
CYTOMEGALOVIRUS ANTIBODIES (IGG, IGM) CYTOMEGALOVIRUS ANTIBODY (IGG)	<0.60	U/mL	U/mL	
	U/mL	Interpretation		
	<0.60	Negative		
	0.60-0.69	Equivocal		
	> or = 0.70	Positive		

A positive result indicates that the patient has antibody to CMV. It does not differentiate between an active or past infection.

CYTOMEGALOVIRUS ANTIBODY (IGM)	<30.00	AU/mL	AU/mL	
	AU/mL	Interpretation		
	<30.00	No Antibody Detected		
	30.00-34.99	Equivocal		
	> or = 35.00	Antibody Detected		

Results from any one IgM assay should not be used as a sole determinant of a current or recent infection. Because an IgM test can yield false positive results and low level IgM antibody may persist for more than 12 months post infection, reliance on a single test result could be misleading. Acute infection is best diagnosed by demonstrating the conversion of IgG from negative to positive. If an acute infection is suspected, consider obtaining a new specimen and submit for both IgG and IgM testing in two or more weeks.

PERFORMING SITE:

SPECIMEN: