

Patient Information	Specimen Information	Client Information

COMMENTS:

Test Name	In Range	Out Of Range	Reference Range	Lab
MEASLES AB (IGG, IGM), DIAGNOSTIC MEASLES AB (IGG), DIAGNOSTIC		24.30 H	AU/mL	

AU/mL	Interpretation
<13.50	Negative
13.50-16.49	Equivocal
>16.49	Positive

The presence of measles IgG suggests immunization or past or current infection with measles virus.

For additional information, please refer to <http://education.QuestDiagnostics.com/faq/FAQ162> (This link is being provided for informational/educational purposes only.)

MEASLES ANTIBODY (IGM)	<1:20	titer
------------------------	-------	-------

Titer	Interpretation
<1:20	Antibody not detected
>or=1:20	Antibody detected

The traditional serologic diagnosis of measles requires a significant rise in antibody titer between acute and convalescent sera. However, detection of IgM antibody in a single specimen may indicate acute disease. Correct interpretation of serologic data depends upon the proper timing of specimen collection in relation to rash onset. This timing is particularly important for interpreting negative IgM results, since IgM antibody peaks approximately 10 days after rash onset and is usually undetectable 30 days after rash onset.

This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics. It has not been cleared or approved by FDA. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes.

For additional information, please refer to <http://education.QuestDiagnostics.com/faq/FAQ162> (This link is being provided for informational/Educational purposes only.)

PERFORMING SITE:

SPECIMEN: