

— Metabolic & Endocrine Health

The endocrine system is the collection of glands, each of which secretes different types of hormones that regulate metabolism, growth and development, tissue function, sexual function, reproduction, sleep and mood, among other things.

The endocrine system is made of eight major glands, which are groups of cells that produce and secrete chemicals. A gland selects and removes materials from the blood, processes them, and secretes the finished chemical product for use somewhere in the body. Almost every organ and cell in the body is affected by the endocrine system.

A group of glands that signal each other in sequence are usually referred to as an axis. One example is the hypothalamic-pituitary-adrenal axis, which coordinates interactions among the hypothalamus, the pituitary gland and the adrenal, also called "suprarenal" glands, which are small, conical organs on top of the kidneys.

The endocrine system sends signals throughout the body, much like the nervous system, but unlike the immediate responses triggered by the nervous system, the effects can take a few hours or weeks. Hormones released from endocrine tissue into the bloodstream where they travel to target tissue to elicit a response.

Endocrine glands are vascular and generally do not have ducts, using intracellular vacuoles, or granules, to store hormones. They differ from, exocrine glands – salivary glands, sweat glands and glands within the gastrointestinal tract – which have ducts or a hollow lumen.


The endocrine system gets some help from organs such as the kidney, liver, heart and gonads, which have secondary endocrine functions. The kidney, for example, secretes hormones such as erythropoietin and renin.

[Hide](#)

Phospholipid Abnormalities

A substance in the body that contains both lipid (fat) and phosphorous; phospholipids are found in all cells throughout the body because they are a major component of the cell membrane, the outermost layer of a cell.

[Hide](#)




B2 GLYCOPROTEIN I (IGM)AB
(U/mL)

<2.0

Range: <20.0

B2 GLYCOPROTEIN I (IGM)AB

(U/mL)



Result Comments

Value	Interpretation
-----	-----
< 20.0	Antibody not detected
> or = 20.0	Antibody detected


B2 GLYCOPROTEIN I (IGA)AB
(U/mL)

<2.0

Range: <20.0

B2 GLYCOPROTEIN I (IGA)AB

(U/mL)



Result Comments

Value	Interpretation
-----	-----
< 20.0	Antibody not detected
> or = 20.0	Antibody detected

The antiphospholipid antibody syndrome (APS) is a clinical-pathologic correlation that includes a clinical event (e.g. arterial or venous thrombosis, pregnancy morbidity) and persistent positive antiphospholipid antibodies (IgM, IgG Cardiolipin or b2GPI antibodies greater than the 99th percentile; or a lupus anticoagulant). International consensus guidelines for APS suggest waiting at least 12 weeks before retesting to confirm antibody persistence. The Systemic Lupus International Collaborating Clinics immunological classification criteria for systemic lupus erythematosus (SLE) include testing for isotype IgA, which has yet to be incorporated into APS criteria. Low level antiphospholipid antibodies may sometimes be detected in the setting of infection, drug therapy or aging.

For additional information, please refer to <http://education.questdiagnostics.com/faq/FAQ109>
(This link is being provided for informational/educational purposes only.)